

Absence of spontaneous magnetic order at non-zero temperature in one- and two-dimensional Heisenberg and XY systems with long-range interactions

P. Bruno*

Max-Planck-Institut für Mikrostrukturphysik, Weinberg 2, D-06120 Halle, Germany[†]

The Mermin-Wagner theorem is strengthened so as to rule out magnetic long-range order at $T > 0$ in one- or two-dimensional Heisenberg and XY systems with *long-range* interactions decreasing as $R^{-\alpha}$ with a sufficiently large exponent α . For *oscillatory* interactions, ferromagnetic long-range order at $T > 0$ is ruled out if $\alpha \geq 1$ ($D = 1$) or $\alpha > 5/2$ ($D = 2$). For systems with *monotonically decreasing* interactions ferro- or antiferromagnetic long-range order at $T > 0$ is ruled out if $\alpha \geq 2D$.

published in: Phys. Rev. Lett. **87**, 137203 (2001)

In a seminal paper exploiting a thermodynamic inequality due to Bogoliubov [1], Mermin and Wagner proved the following important theorem [2]: “For one- or two-dimensional Heisenberg systems with isotropic interactions and such that the interactions are short-ranged, namely which satisfy the condition

$$\sum_{\mathbf{R}} \mathbf{R}^2 |J_{\mathbf{R}}| < +\infty, \quad (1)$$

there can be no spontaneous ferro- or antiferromagnetic long range order at $T > 0$.” [3] In view of the fact that most of our knowledge on critical phenomena and phase transitions is based upon approximate theories, the few known rigorous results [4] such as the Mermin-Wagner theorem are of key importance, for they allow to test the validity of approximate theories.

For interactions with a finite range or with an exponential decay, the condition (1) is trivially satisfied. For interactions with a power-law decay $|J_{\mathbf{R}}| \propto R^{-\alpha}$, condition (1) is satisfied provided that $\alpha > D + 2$, where D is the dimensionality. In metallic magnetic systems the exchange interactions are of the Ruderman-Kittel-Kasuya-Yosida (RKKY) type [5], which have a long range oscillatory behavior for large R : $J_{\mathbf{R}} \propto R^{-D} \cos(q_0 R + \phi)$. It is obvious that the RKKY interactions do not satisfy the criterion of short-rangedness (1), so that no conclusion on the magnetism one- and two-dimensional RKKY systems can be obtained from the Mermin-Wagner theorem. This situation is highly unsatisfactory, in view of the fact that most magnetic ultrathin films investigated experimentally consist of metals and alloys.

In the present paper, I extend the result of Mermin and Wagner to Heisenberg and XY systems with a *long range interaction*, i.e., which do not satisfy the condition (1), with particular emphasis on systems with oscillatory interactions. Results for systems with monotonically decaying interactions are also presented. More specifically, I consider one- and two-dimensional systems described by the following Hamiltonian:

$$H = - \sum_{\mathbf{R}, \mathbf{R}'} [J'_{\mathbf{R}-\mathbf{R}'} S_{\mathbf{R}}^x S_{\mathbf{R}'}^x + J_{\mathbf{R}-\mathbf{R}'} (S_{\mathbf{R}}^y S_{\mathbf{R}'}^y + S_{\mathbf{R}}^z S_{\mathbf{R}'}^z)] + K \sum_{\mathbf{R}} (S_{\mathbf{R}}^x)^2 - B \sum_{\mathbf{R}} S_{\mathbf{R}}^z. \quad (2)$$

We also define the Fourier transform of the spin operator $\mathbf{S}_{\mathbf{R}}$, $\mathbf{S}(\mathbf{k}) \equiv \sum_{\mathbf{R}} \mathbf{S}_{\mathbf{R}} e^{-i\mathbf{k} \cdot \mathbf{R}}$, as well as $E(\mathbf{k}) \equiv \sum_{\mathbf{R}} J_{\mathbf{R}} (1 - e^{i\mathbf{k} \cdot \mathbf{R}}) = E(-\mathbf{k})$, and $\tilde{E}(\mathbf{k}) \equiv \sum_{\mathbf{R}} |J_{\mathbf{R}}| (1 - e^{i\mathbf{k} \cdot \mathbf{R}}) = \tilde{E}(-\mathbf{k})$. For $K = 0$ and $J'_{\mathbf{R}} = J_{\mathbf{R}}$ the above Hamiltonian corresponds to the isotropic Heisenberg model. In the general case, it corresponds to a system with uniaxial anisotropy of axis x , with a single-spin anisotropy (for $S \geq 1$) and/or a two-spin exchange anisotropy. Depending value of the anisotropy parameters K and $J'_{\mathbf{R}} - J_{\mathbf{R}}$, one therefore has either an isotropic system, an XY-like system with yz easy plane, or an Ising-like model with x easy axis (in the latter case, one merely shows that the spontaneous magnetization has to be along the x axis). For such systems (with arbitrary values of parameters $J'_{\mathbf{R}}$ and K), I prove the following results:

Theorem 1: “A D -dimensional ($D = 1$ or 2) Heisenberg or XY system for which one can find some numbers $k_0 > 0$ (with k_0 belonging to the first BZ), $\alpha > 0$, $\beta \leq 1$ such that for $|\mathbf{k}| < k_0$ and $|\mathbf{k}'| < k_0$

$$|E(\mathbf{k})| \leq \alpha |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta, \quad (3a)$$

$$|E(\mathbf{k}' - \mathbf{k}) + E(\mathbf{k}' + \mathbf{k}) - 2E(\mathbf{k}')| \leq \alpha |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta, \quad (3b)$$

$$\frac{1}{A_D} \int d^D \mathbf{k}'' |E(\mathbf{k}'' - \mathbf{k}) + E(\mathbf{k}'' + \mathbf{k}) - 2E(\mathbf{k}'')| \leq \alpha |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta, \quad (3c)$$

cannot be ferromagnetic.” [6]

Corollary 1: “A D -dimensional ($D = 1$ or 2) Heisenberg or XY system with long-range oscillatory interactions of the form

$$J_{\mathbf{R}} \propto \frac{\cos(q_0 R + \phi)}{R^\alpha}, \quad (4)$$

with $q_0 \neq 0$ belonging to the first Brillouin zone (BZ) and $\alpha \geq 1$ ($D = 1$), or $\alpha > 3/2$ ($D = 2$, $\sin(\phi + \pi/4) = 0$), or $\alpha > 5/2$ ($D = 2$, $\sin(\phi + \pi/4) \neq 0$) cannot be ferromagnetic.”

Theorem 2: “A D -dimensional ($D = 1$ or 2) Heisenberg or XY system satisfying

$$\frac{1}{A_D} \int d^D \mathbf{k} \frac{1}{\tilde{E}(\mathbf{k})} = +\infty, \quad (5)$$

where A_D is the measure of the first Brillouin zone (BZ), cannot be ferro- or antiferromagnetic.”

Corollary 2: “A D -dimensional ($D = 1$ or 2) Heisenberg or XY system with interactions monotonically decaying as $|J_{\mathbf{R}}| \propto R^{-\alpha}$ with $\alpha \geq 2D$ cannot be ferro- or antiferromagnetic.”

Proof of Theorem 1: Like for the Mermin-Wagner theorem, the proof of Theorem 1 relies on using the Bogoliubov inequality to prove that the z component of the spontaneous magnetization $\lim_{B \rightarrow 0} |\langle S^z \rangle|$ vanishes for any finite temperature. The proof of Theorem 1 is however significantly more delicate than Mermin-Wagner’s one.

Our proof proceeds by *reductio ad absurdo*. Let us assume that the system is ferromagnetic. This implies that one can find a temperature $T_0 > 0$ and a quantity $m_0 > 0$ such that for any temperature $T \leq T_0$, one has $|\langle S^z \rangle| \geq m_0$.

The Bogoliubov inequality [1] states that $\langle \{A, A^\dagger\} \rangle \langle [[C, H], C^\dagger] \rangle \geq 2k_B T |\langle [C, A] \rangle|^2$, where A and C are arbitrary operators, $[A, B]$ is the commutator of A and B , $\{A, B\}$ is the anticommutator of A and B , and $\langle A \rangle$ is the thermodynamic average of A . The two factors on the left-hand side of the inequality are ≥ 0 . We use Bogoliubov’s inequality for $A \equiv S^y(-\mathbf{k})$ and $C \equiv S^x(\mathbf{k})$. Straightforward algebra together with the inequality $\langle (S_{\mathbf{R}}^y)^2 \rangle \leq S^2$ then yield the following inequality:

$$S^2 \geq \frac{1}{A_D} \int d^D \mathbf{k} \frac{k_B T \langle S^z \rangle^2}{B \langle S^z \rangle + 2 \sum_{\mathbf{R}} J_{\mathbf{R}} [1 - \cos(\mathbf{k} \cdot \mathbf{R})] \langle S_0^y S_{\mathbf{R}}^y + S_0^z S_{\mathbf{R}}^z \rangle}. \quad (6)$$

Note that the above result is independent of $J'_{\mathbf{R}}$ and K , and depends only on $J_{\mathbf{R}}$. So far, our proof follows exactly the one of Mermin and Wagner. From this point, Mermin and Wagner proceed by stating that if the condition (1) is satisfied, the (positive) denominator $\Delta(\mathbf{k})$ of the above equation can be majorated near $\mathbf{k} = 0$ by an expression of the form $|B|S + \alpha k^2$ (with $\alpha > 0$), from which they then easily that the spontaneous magnetization vanishes.

In order to obtain a stronger theorem, i.e., to rule out

ferromagnetism for a broader class of systems, one therefore needs to find a tighter majoration of $\Delta(\mathbf{k})$ than used by Mermin and Wagner. One can check easily that a majoration of the form $\Delta(\mathbf{k}) \leq |B|S + \gamma |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta$ near $\mathbf{k} = 0$, with $\gamma > 0$ and $\beta \leq 1$, would be sufficient for our purpose.

To this aim, we rewrite the denominator $\Delta(\mathbf{k})$ of Eq. (6) as

$$\Delta(\mathbf{k}) = B \langle S^z \rangle + 2 \langle S^z \rangle^2 E(\mathbf{k}) + \frac{1}{A_D} \int d^D \mathbf{k}' [E(\mathbf{k}' - \mathbf{k}) + E(\mathbf{k}' + \mathbf{k}) - 2E(\mathbf{k}')] F(\mathbf{k}'), \quad (7)$$

with

$$\begin{aligned} F(\mathbf{k}) &\equiv \sum_{\mathbf{R}} e^{-i\mathbf{k} \cdot \mathbf{R}} \langle S_0^y S_{\mathbf{R}}^y + \delta S_0^z \delta S_{\mathbf{R}}^z \rangle \\ &= \frac{1}{N} \langle |S^y(\mathbf{k})|^2 + |\delta S^z(\mathbf{k})|^2 \rangle \\ &= k_B T [\chi^{yy}(\mathbf{k}) + \chi^{zz}(\mathbf{k})] \end{aligned} \quad (8)$$

where we have introduced the longitudinal fluctuation $\delta S_{\mathbf{R}}^z \equiv S_{\mathbf{R}}^z - \langle S^z \rangle$ and the transverse and longitudinal $\mathbf{k}$ -dependent susceptibilities $\chi^{yy}(\mathbf{k})$ and $\chi^{zz}(\mathbf{k})$, respectively. Since we are assuming that the system has a long-range ferromagnetic order, one can argue that $F(\mathbf{k})$ has to be finite for $\mathbf{k} \neq 0$, otherwise the ferromagnetic or-

der would be unstable against fluctuations of non-zero wavevector. On the other hand $F(\mathbf{k})$ certainly diverges for $\mathbf{k} = 0$ due to the divergence of the static transverse susceptibility of systems having a continuous rotational invariance. Nevertheless, one has

$$\frac{1}{A_D} \int d^D \mathbf{k} F(\mathbf{k}) = \langle (S^y)^2 + (S^z)^2 \rangle \leq S(S+1). \quad (9)$$

In view of the above remarks concerning $F(\mathbf{k})$, there exists a number $F_0 > 0$ such that $0 \leq F(\mathbf{k}) \leq F_0$ for $|\mathbf{k}| > k_0$. Making use of conditions (3b,c), one can therefore write, for $|\mathbf{k}| < k_0$:

$$\begin{aligned}
A(\mathbf{k}) &\equiv \frac{1}{A_D} \int d^D \mathbf{k}' |E(\mathbf{k}' - \mathbf{k}) + E(\mathbf{k}' + \mathbf{k}) - 2E(\mathbf{k}')| F(\mathbf{k}'), \\
&\leq \frac{1}{A_D} \int_{|\mathbf{k}'| < k_0} d^D \mathbf{k}' \alpha |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta F(\mathbf{k}') + \frac{1}{A_D} \int_{|\mathbf{k}'| > k_0} d^D \mathbf{k}' |E(\mathbf{k}' - \mathbf{k}) + E(\mathbf{k}' + \mathbf{k}) - 2E(\mathbf{k}')| F_0 \\
&\leq \frac{1}{A_D} \int d^D \mathbf{k}' \alpha |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta F(\mathbf{k}') + \frac{1}{A_D} \int d^D \mathbf{k}' |E(\mathbf{k}' - \mathbf{k}) + E(\mathbf{k}' + \mathbf{k}) - 2E(\mathbf{k}')| F_0 \\
&\leq [S(S+1) + F_0] \alpha |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta.
\end{aligned} \tag{10}$$

Combining this result with condition (3a), we obtain that

$$0 \leq \Delta(\mathbf{k}) \leq |B|S + \gamma |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta, \tag{11}$$

for $|\mathbf{k}| < k_0$, with $\gamma \equiv S^2 + S(S+1) + F_0 > 0$. Combining the above result with inequality (6), one obtains

$$S^2 \geq \frac{1}{A_D} \int_{|\mathbf{k}'| < k_0} d^D \mathbf{k}' \frac{k_B T \langle S^z \rangle^2}{|B|S + \gamma |\mathbf{k}|^D |\ln(|\mathbf{k}|^{-1})|^\beta}, \tag{12}$$

from which one shows immediately that $\lim_{B \rightarrow 0} \langle S^z \rangle = 0$ for $T > 0$, which is in contradiction with our hypothesis that the system is ferromagnetic. This completes the proof of Theorem 1 by *reductio ad absurdum*. $\square$

Proof of Corollary 1: Consider interaction of the form Eq. (4). To investigate the behavior of $E(\mathbf{k})$ in the vicinity of $\mathbf{k} = 0$, we can substitute the discrete sum over $\mathbf{R}$ by an integral. One can show easily that, for any value of α , $E(\mathbf{k})$ is analytical for all wavevectors except for $|\mathbf{k}| = q_0$. Therefore, for $\mathbf{k} \rightarrow 0$,

$$E(\mathbf{k}) \propto |\mathbf{k}|^2 \quad (\forall \alpha), \tag{13}$$

and, for $\mathbf{k} \rightarrow 0$ and $|\mathbf{k}| < |\mathbf{k}'| - q_0$,

$$|E(\mathbf{k}' - \mathbf{k}) + E(\mathbf{k}' + \mathbf{k}) - 2E(\mathbf{k}')| \propto |\mathbf{k}|^2 \quad (\forall \alpha). \tag{14}$$

The integral over $\mathbf{k}''$ in (3c) is dominated by the non-analyticity of $E(\mathbf{k})$ in the vicinity of $|\mathbf{k}| = q_0$. Let us therefore study $\varepsilon(\kappa)$, the singular part of $E(\mathbf{k})$ for $\kappa \equiv |\mathbf{k}| - q_0 \rightarrow 0$. We get

$$\varepsilon(\kappa) \propto \begin{cases} [A + B \operatorname{sgn}(\kappa)] |\kappa|^{\alpha-1} & (\alpha < 1), \\ A \ln |\kappa|^{-1} + B \operatorname{sgn}(\kappa) & (\alpha = 1), \\ B \operatorname{sgn}(\kappa) |\kappa|^{\alpha-1} & (1 < \alpha < 2), \\ B \operatorname{sgn}(\kappa) |\kappa| \ln |\kappa|^{-1} & (\alpha = 2), \\ B \operatorname{sgn}(\kappa) |\kappa| & (\alpha > 2), \end{cases} \tag{15}$$

for $D = 1$, with $A \propto \cos \phi$ and $B \propto \sin \phi$, and

$$\varepsilon(\kappa) \propto \begin{cases} [A + B \operatorname{sgn}(\kappa)] |\kappa|^{\alpha-3/2} & (\alpha < 3/2), \\ A \ln |\kappa|^{-1} + B \operatorname{sgn}(\kappa) & (\alpha = 3/2), \\ B \operatorname{sgn}(\kappa) |\kappa|^{\alpha-3/2} & (3/2 < \alpha < 5/2), \\ B \operatorname{sgn}(\kappa) |\kappa| \ln |\kappa|^{-1} & (\alpha = 5/2), \\ B \operatorname{sgn}(\kappa) |\kappa| & (\alpha > 5/2), \end{cases} \tag{16}$$

for $D = 2$, with $A \propto \cos(\phi + \pi/4)$ and $B \propto \sin(\phi + \pi/4)$. Let $C(\mathbf{k})$ be the left-hand side expression in Eq. (3c). We then get, for $\mathbf{k} \rightarrow 0$,

$$C(k) \propto \begin{cases} +\infty & (\alpha \leq 0), \\ |k|^\alpha & (0 < \alpha < 1), \\ |k| \ln |k|^{-1} & (\alpha = 1, \cos \phi \neq 0), \\ |k| & (\alpha = 1, \cos \phi = 0), \\ |k|^2 & (\alpha > 1, \sin \phi = 0), \\ |k| & (1 < \alpha < 2, \sin \phi \neq 0), \\ |k|^2 \ln |k|^{-1} & (\alpha = 2, \sin \phi \neq 0), \\ |k|^2 & (\alpha > 2, \sin \phi \neq 0), \end{cases} \tag{17}$$

for $D = 1$, and

$$C(\mathbf{k}) \propto \begin{cases} +\infty & (\alpha \leq 1/2), \\ |\mathbf{k}|^{\alpha-1/2} & (1/2 < \alpha < 3/2), \\ |\mathbf{k}| \ln |\mathbf{k}|^{-1} & (\alpha = 3/2, \cos(\phi + \pi/4) \neq 0), \\ |\mathbf{k}| & (\alpha = 3/2, \cos(\phi + \pi/4) = 0), \\ |\mathbf{k}|^2 & (\alpha > 3/2, \sin(\phi + \pi/4) = 0), \\ |\mathbf{k}|^{\alpha-1/2} & (3/2 < \alpha < 5/2, \sin(\phi + \pi/4) \neq 0), \\ |\mathbf{k}| \ln |\mathbf{k}|^{-1} & (\alpha = 5/2, \sin(\phi + \pi/4) \neq 0), \\ |\mathbf{k}|^2 & (\alpha > 5/2, \sin(\phi + \pi/4) \neq 0), \end{cases} \tag{18}$$

for $D = 2$. From Eqs. (13,14), it follows that conditions (3a,b) are satisfied for all values of α . From Eq. (17,18), one gets that condition (3c) is fulfilled for $\alpha \geq 1$ ($D = 1$), and for $\alpha > 3/2$ ($D = 2$, $\sin(\phi + \pi/4) = 0$) or $\alpha > 5/2$ ($D = 2$, $\sin(\phi + \pi/4) \neq 0$), which completes the proof of Corollary 1. $\square$

Proof of Theorem 2: Theorem 2 is an immediate generalization of the Mermin-Wagner theorem. For the sake of simplicity, we detail the proof only for the case of ferromagnetism; the extension to the case of antiferromagnetism is immediate by introducing a staggered field and staggered magnetization as done by Mermin and Wagner.

The proof follows the one of Theorem 1 until Eq. (6). Then, by using $|\langle S_0^y S_{\mathbf{R}}^y + S_0^z S_{\mathbf{R}}^z \rangle| \leq S(S+1)$:

$$\Delta(\mathbf{k}) \leq |B|S + 2S(S+1)\tilde{E}(\mathbf{k}), \tag{19}$$

From condition (5) it follows immediately that $\lim_{B \rightarrow 0} |\langle S^z \rangle| = 0$ for $T > 0$, which is in contradiction with our hypothesis that the system is ferromagnetic. This completes the proof of Theorem 2 by *reductio ad absurdo*. $\square$

Proof of Corollary 2: Let us consider the behavior of $\tilde{E}(\mathbf{k})$ in the vicinity of $\mathbf{k} = 0$ for $|J_{\mathbf{R}}| \propto R^{-\alpha}$. In this regime, the discrete sum over $\mathbf{R}$ can be replaced by an integral, and one shows easily that, for $|\mathbf{k}| \rightarrow 0$,

$$\tilde{E}(\mathbf{k}) \propto \begin{cases} |k|^{\alpha-D} & (D < \alpha < D+2), \\ |k|^2 \ln(|k|^{-1}) & (\alpha = D+2), \\ |k|^2 & (\alpha > D+2). \end{cases} \quad (20)$$

It then follows immediately that condition (5) is satisfied if $\alpha \geq 2D$, which completes the proof of Corollary 2. $\square$

As for the Mermin-Wagner theorem, the above theorems and corollaries can be extended also to one-dimensional systems of arbitrary finite cross-section and to two-dimensional systems of arbitrary finite thickness.

The importance of Theorem 1 and Corollary 1 is that the class of systems for which ferromagnetism is rigorously ruled out is significantly increased. In particular, we obtain interesting results for systems with RKKY interactions ($\alpha = D$). For one-dimensional RKKY systems, ferromagnetism is rigorously ruled out. For two-dimensional RKKY systems, the results are less satisfactory, since ferromagnetism can be strictly ruled out only for the particular case with $\sin(\phi + \pi/4) = 0$. For the general case ($\sin(\phi + \pi/4) \neq 0$), it does not seem possible to rule out ferromagnetism for $\alpha = D = 2$ without analyzing in detail the behavior of $F(\mathbf{k})$ near $|\mathbf{k}| = q_0$. If one could prove that $F(\mathbf{k})$ has no singularity more singular than a square-root singularity at $|\mathbf{k}| = q_0$, then the absence of ferromagnetism would follow from Eq. (18). This situation is somehow puzzling. Indeed, within the (non rigorous) linearized spin-wave theory, as well as the random phase approximation (RPA) Green's function method, having $E(\mathbf{k}) \propto k^2$ is sufficient to rule out ferromagnetism in two dimensions, which implies that these theories rule out ferromagnetism in systems with oscillatory interactions, for any value of α . In view of these considerations, I therefore propose the following result:

Conjecture 1: “A D -dimensional ($D = 1$ or 2) Heisenberg or XY system for which there exists some finite $\mathbf{k}$ region with $E(\mathbf{k}) < 0$, or satisfying

$$\frac{1}{A_D} \int d^D \mathbf{k} \frac{1}{E(\mathbf{k})} = +\infty, \quad (21)$$

cannot be ferromagnetic.”

From Eq. (13), it then follows immediately:

Corollary 3: “If Conjecture 1 is true, then a D -dimensional ($D = 1$ or 2) Heisenberg or XY system with

long-range oscillatory interactions of the form (4) cannot be ferromagnetic, for any value of α .”

Let us now discuss the results of Theorem 2 and Corollary 2. They essentially constitute an extension to XY systems of a result obtained earlier by Joyce [7]. For the one-dimensional (classical) Heisenberg and XY systems with positive (ferromagnetic) long range interaction, Fröhlich *et al.* proved the existence of ferromagnetism if $1 < \alpha < 2$ [8]. This is to be contrasted with one-dimensional Ising with long range interaction, for which ferromagnetism is ruled out if $\alpha > 2$ [9], whereas ferromagnetism exists for $1 < \alpha \leq 2$ [10]. The Heisenberg and XY systems therefore only differ from the Ising system only on the borderline $\alpha = 2$: there ferromagnetism is excluded for the Heisenberg and XY cases, but exists for the Ising case, illustrating the stronger tendency towards ordering displayed by the Ising model.

I wish to express my thanks to V. Dugaev and G. Bouzerar for helpful discussions.

* Electronic address: bruno@mpi-halle.de

† <http://www.mpi-halle.de>

- [1] N.N. Bogoliubov, Physik Abhandl. Sowjetunion **6**, 1 (1962).
- [2] N.D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1133 (1966).
- [3] For the sake of concision, throughout this paper, a system will be termed “ferromagnetic” (respectively “antiferromagnetic”) (or not) without further precision if it possesses (or not) “ferromagnetic (respectively “antiferromagnetic”) long range order below some non-zero Curie temperature”, without prejudice of the magnetic order exhibited in the ground state.
- [4] D. Ruelle, *Statistical Mechanics: Rigorous Results*, Benjamin (1969); R.B. Griffiths, in *Phase Transitions and Critical Phenomena*, vol. 1, ed. by C. Domb and M.S. Green, Academic Press (New York, 1972), p. 7; B. Simon, *The Statistical Mechanics of Lattice Gases*, vol. 1, Princeton University Press (Princeton, 1993).
- [5] M.A. Ruderman and C. Kittel, Phys. Rev. **96**, 99 (1954); T. Kasuya, Prog. Theor. Phys. **16**, 45 (1956); K. Yosida, Phys. Rev. **106**, 893 (1957).
- [6] The continuous integration over wavevectors indicates that we are taking the thermodynamic limit. Throughout the paper, all integrations over wavevectors are strictly restricted to the first BZ, unless explicitly further restricted.
- [7] G.S. Joyce, J. Phys. C: Solid State Phys. **2**, 1531 (1969).
- [8] J. Fröhlich, R. Israel, E.H. Lieb, and B. Simon, Commun. Math. Phys. **62**, 1 (1978).
- [9] D. Ruelle, Commun. Math. Phys. **9**, 267 (1968).
- [10] F.J. Dyson, Commun. Math. Phys. **12**, 91 (1969); J. Fröhlich and T. Spencer, Commun. Math. Phys. **84**, 87 (1982).